LISTING OF THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

- 1. (**Previously Presented**) A method for preventing infection of a subject by a microbe comprising: administering a compound that downmodulates the expression or activity of a microbial transcription factor to a subject at risk of developing an infection, wherein said downmodulation of the microbial transcription factor reduces virulence of the microbe, such that infection of the subject is prevented.
- 2. **(Original)** The method of claim 1, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.
- 3. (**Original**) The method of claim 1, wherein the transcription factor is a member of the MarA family of transcription factors.
- 4. **(Previously Presented)** The method of claim 1, further comprising administering an antibiotic.
- 5. (**Previously Presented**) A method for preventing urinary tract infection of a subject by a microbe comprising: administering a compound that downmodulates the expression or activity of a microbial transcription factor to a subject at risk of developing a urinary tract infection such that infection of the subject is prevented.
- 6. (**Previously Presented**) A method for preventing prostatitis in a subject by a microbe comprising: administering a compound that downmodulates the expression or activity of a microbial transcription factor to a subject at risk of developing prostatitis such that infection of the subject is prevented.
- 7. (Withdrawn) A method for reducing virulence of a microbe comprising: administering a compound that modulates the expression or activity of a microbial transcription factor to a subject at risk of developing an infection with the microbe such that virulence of the microbe is reduced.

8. **(Withdrawn)** The method of claim 7, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.

- 9. (Withdrawn) The method of claim 7, wherein the transcription factor is a member of the MarA family of transcription factors.
- 10. (Withdrawn) The method of claim 7, further comprising administering an antibiotic.
- 11. (Withdrawn) A method for treating a microbial infection in a subject comprising: administering a compound that modulates the expression or activity of a microbial transcription factor to a subject having a microbial infection, wherein said modulation of the microbial transcription factor reduces virulence of the microbe, such that infection of the subject is treated.
- 12. **(Withdrawn)** The method of claim 11, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.
- 13. (Withdrawn) The method of claim 11, wherein the transcription factor is a member of the MarA family of transcription factors.
- 14. (Withdrawn) The method of claim 11, further comprising administering an antibiotic.
- 15. (Withdrawn) A method for treating a microbial urinary tract infection in a subject comprising: administering a compound that modulates the expression or activity of a microbial transcription factor to a subject having a microbial urinary tract infection such that infection of the subject is treated.
- 16. (Withdrawn) A method for treating prostatitis in a subject comprising: administering a compound that modulates the expression or activity of a transcription factor to a subject having prostatitis such that infection of the subject is treated.
- 17. **(Withdrawn)** The method of claim 15, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.
- 18. (Withdrawn) The method of claim 15, wherein the transcription factor is a member of the MarA family of transcription factors.

19. (Withdrawn) The method of claim 15, further comprising administering an antibiotic.

- 20. (Withdrawn) A method for reducing virulence in a microbe comprising: administering a compound that inhibits the expression or activity of a transcription factor to a subject having a microbial infection such that virulence of the microbe is reduced.
- 21. (Withdrawn) The method of claim 20, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.
- 22. (Withdrawn) The method of claim 20, wherein the transcription factor is a member of the MarA family of transcription factors.
- 23. (Withdrawn) The method of claim 20, further comprising administering an antibiotic.
- 24. (Withdrawn) A method for evaluating the effectiveness of a compound that modulates the expression or activity of a microbial transcription factor at inhibiting microbial virulence comprising: infecting a non-human animal with a microbe, wherein the ability of the microbe to establish an infection in the non-human animal requires that the microbe colonize the animal; administering the compound that modulates the expression or activity of the microbial transcription factor to the non-human animal; and determining the level of infection of the non-human animal, wherein the ability of the compound to reduce the level of infection of the animal indicates that the compound is effective at inhibiting microbial virulence.
- 25. (Withdrawn) The method of claim 24, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.
- 26. (Withdrawn) The method of claim 24, wherein the transcription factor is a member of the MarA family of transcription factors.
- 27. (Withdrawn) The method of claim 24, further comprising administering an antibiotic.
- 28. (Withdrawn) The method of claim 24, wherein the level of infection of the non-human animal is determined by measuring the ability of the microbe to colonize the tissue of the non-human animal.

29. (Withdrawn) The method of claim 24, wherein the level of infection of the non-human animal is determined by enumerating the number of microbes present in the tissue of the non-human animal.

- 30. (Withdrawn) A method for identifying a compound for treating microbial infection, comprising: innoculating a non-human animal with a microbe, wherein the ability of the microbe to establish an infection in the non-human animal requires that the microbe colonize the animal; administering a compound which reduces the expression or activity of a microbial transcription factor to the animal, and determining the effect of the test compound on the ability of the microbe to colonize the animal, such that a compound for treating microbial infection is identified.
- 31. **(Withdrawn)** The method of claim 30, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.
- 32. (Withdrawn) The method of claim 30, wherein the transcription factor is a member of the MarA family of transcription factors.
- 33. (Withdrawn) The method of claim 30, wherein the level of infection of the non-human animal is determined by measuring the ability of the microbe to colonize the tissue of the non-human animal.
- 34. (Withdrawn) The method of claim 30, wherein the level of infection of the non-human animal is determined by enumerating the number of microbes present in the tissue of the non-human animal.
- 35. (Withdrawn) A method for identifying a compound for reducing microbial virulence, comprising: inoculating a non-human animal with a microbe, wherein the ability of the microbe to establish an infection in the non-human animal requires that the microbe colonize the animal; administering a compound which reduces the expression or activity of a microbial transcription factor to the animal, and determining the effect of the test compound on the ability of the microbe to colonize the animal, such that a compound for reducing microbial virulence is identified.
- 36. (Withdrawn) The method of claim 35, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.

37. (Withdrawn) The method of claim 35, wherein the transcription factor is a member of the MarA family of transcription factors.

- 38. (Withdrawn) The method of claim 35, wherein the level of infection of the non-human animal is determined by measuring the ability of the microbe to colonize the tissue of the non-human animal.
- 39. (Withdrawn) The method of claim 35, wherein the level of infection of the non-human animal is determined by enumerating the number of microbes present in the tissue of the non-human animal.
- 40. (Withdrawn) A method for identifying transcription factors which promote microbial virulence comprising: creating a microbe in which a transcription factor to be tested is misexpressed; introducing the microbe into a non-human animal; wherein the ability of the microbe to establish an infection in the non-human animal requires that the microbe colonize the animal; and determining the ability of the microbe to colonize the animal, wherein a reduced ability of the microbe to colonize the animal as compared to a wild-type microbial cell identifies the transcription factor as a transcription factor which promotes microbial virulence.
- 41. **(Withdrawn)** The method of claim 40, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.
- 42. (Withdrawn) The method of claim 40, wherein the transcription factor is a member of the MarA family of transcription factors.
- 43. (Withdrawn) The method of claim 40, wherein the level of infection of the non-human animal is determined by measuring the ability of the microbe to colonize the tissue of the non-human animal.
- 44. (Withdrawn) The method of claim 40, wherein the level of infection of the non-human animal is determined by enumerating the number of microbes present in the tissue of the non-human animal.
- 45. (Withdrawn) A method for reducing the ability of a microbe to adhere to an abiotic surface comprising: contacting the abiotic surface or the microbe with a compound that

modulates the activity of a transcription factor such that the ability of the microbe to adhere to the abiotic surface is reduced.

- 46. (Withdrawn) The method of claim 45, wherein the transcription factor is a member of the AraC-XylS family of transcription factors.
- 47. (Withdrawn) The method of claim 45, wherein the transcription factor is a member of the MarA family of transcription factors.
- 48. (Withdrawn) The method of claim 45, further comprising contacting the abiotic surface or the microbe with a second agent that is effective at controlling the growth of the microbe.
- 49. (Withdrawn) The method of claim 45, wherein the abiotic surface is selected from the group consisting of: stents, catheters, and prosthetic devices.
- 50. (Withdrawn) A pharmaceutical composition comprising a compound that modulates the activity or expression of a microbial transcription factor and a pharmaceutically acceptable carrier, wherein the compound reduces microbial virulence.
- 51. (Withdrawn) A pharmaceutical composition comprising a compound that modulates the activity or expression of a microbial transcription factor and an antibiotic in a pharmaceutically acceptable carrier.
- 52. (**Previously Presented**) A method for treating or preventing infection of a subject by a microbe comprising: administering a compound that downmodulates the expression or activity of a microbial transcription factor to a subject at risk of developing an infection, wherein said downmodulation of the microbial transcription factor reduces virulence of the microbe, such that infection of the subject is treated or prevented.